

University of Dundee

## The specificities of small molecule inhibitors of the TGF beta and BMP pathways

Vogt, Janis; Traynor, Ryan; Sapkota, Gopal P.

*Published in:*  
Cellular Signalling

*DOI:*  
[10.1016/j.cellsig.2011.06.019](https://doi.org/10.1016/j.cellsig.2011.06.019)

*Publication date:*  
2011

*Document Version*  
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

### *Citation for published version (APA):*

Vogt, J., Traynor, R., & Sapkota, G. P. (2011). The specificities of small molecule inhibitors of the TGF beta and BMP pathways. *Cellular Signalling*, 23(11), 1831-1842. <https://doi.org/10.1016/j.cellsig.2011.06.019>

### General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

THE SPECIFICITIES OF SMALL MOLECULE INHIBITORS OF  
THE TGFβ AND BMP PATHWAYS

Janis Vogt<sup>1</sup>, Ryan Traynor<sup>2</sup>, Gopal P Sapkota<sup>1</sup>

<sup>1</sup>MRC Protein Phosphorylation Unit, College of Life Sciences, <sup>2</sup> Division of Signal Transduction Therapy (DSTT), University of Dundee, Dow Street, Dundee, DD1 5EH, Scotland, UK.  
Address correspondence to: Dr. Gopal Sapkota, Sir James Black Centre, Dow Street, Dundee, DD1 5EH, Scotland, UK; Phone: ++44 (0)1382 386330; Fax: ++44 (0)1382 223778; E-mail: [g.sapkota@dundee.ac.uk](mailto:g.sapkota@dundee.ac.uk)

## Abstract

Small molecule inhibitors of type 1 receptor serine threonine kinases (ALKs1-7), the mediators of TGF $\beta$  and BMP signals, have been employed extensively to assess their physiological roles in cells and organisms. While all of these inhibitors have been reported as “selective” inhibitors of specific ALKs, extensive specificity tests against a wide array of protein kinases have not been performed. In this study, we examine the specificities and potencies of the most frequently used small molecule inhibitors of the TGF $\beta$  pathway (SB-431542, SB-505124, LY-364947 and A-83-01) and the BMP pathway (Dorsomorphin and LDN-193189) against a panel of up to 123 protein kinases covering a broad spectrum of the human kinome. We demonstrate that the inhibitors of the TGF $\beta$  pathway are relatively more selective than the inhibitors of the BMP pathway. Based on our specificity and potency profile and published data, we recommend SB-505124 as the most suitable molecule for use as an inhibitor of ALKs 4, 5 & 7 and the TGF $\beta$  pathway. We do not recommend Dorsomorphin, also called Compound C, for use as an inhibitor of the BMP pathway. Although LDN-193189, a Dorsomorphin derivative, is a very potent inhibitor of ALK2/3 and the BMP-pathway, we found that it potently inhibited a number of other protein kinases at concentrations sufficient to inhibit ALK2/3 and its use as a selective BMP-pathway inhibitor has to be considered cautiously. Our observations have highlighted the need for caution when using these small molecule inhibitors to assess the physiological roles of BMP and TGF $\beta$  pathways.

**Keywords:** TGF-beta, BMP, Kinase, Inhibitors, LDN-193189, SB-505124

**Abbreviations:** ActR-IIA, activin A receptor, type IIA; ActR-IIB, activin A receptor, type IIB; ALK, Activin receptor-like kinase; AMH, anti-müllerian hormone; AMHR-II, anti-müllerian hormone receptor type II; BMP, bone morphogenetic protein; BMPR-II, bone morphogenetic protein receptor type II; GDF, growth differentiation factor; TGF $\beta$ , transforming growth factor- $\beta$ ; TGF $\beta$ R-II, transforming growth factor- $\beta$  receptor type II.

## 1. Introduction

The signalling pathways downstream of the transforming growth factor beta (TGF $\beta$ ) family of cytokines, that comprise some 42 members, control plethora of cellular processes including proliferation, differentiation, extra-cellular matrix production, motility, survival and fate [1]. Aberrant TGF $\beta$  signalling pathways are associated with many human diseases, including bone diseases, immune-suppression, fibrosis, cancer progression and metastasis [2-7]. Hence targeted disruption of specific TGF $\beta$  signalling components by small molecules or other means provides potential therapeutic opportunities. Over the past few years, BMP/TGF $\beta$  type I and type II receptor serine threonine protein kinases, the transducers of BMP and TGF $\beta$  signals, have been targeted for development of small molecule inhibitors. Specific small molecule inhibitors of these protein kinases not only provide a flexible, rapid and cost-effective means of inhibiting their targets in cells and tissues but also potentially could have many therapeutic applications.

The TGF $\beta$  family of ligands is broadly divided into two groups based on their ability to trigger the activation of specific Smad transcription factors, the intracellular mediators of TGF $\beta$  signals. The TGF $\beta$  subfamily (which includes TGF $\beta$ , Activin and Nodal) activates Smads 2 and 3, while the BMP subfamily (which includes BMPs, GDFs and AMH) activates Smads 1, 5 and 8 [1]. The ligands exist as homo- or hetero-dimers and bind to specific sets of type II and type I receptors, which are serine-threonine protein kinases, and thus result in a large ligand-receptor complex involving a ligand dimer, two type II and two type I receptor molecules [1, 8]. The formation of ligand-receptor complex facilitates the constitutively active type II receptor kinases to phosphorylate and activate the type I receptor kinases [1, 9]. In all, there are five type II receptors (ActR-IIA, ActR-IIB, BMPR-II, AMHR-II and TGF $\beta$ R-II) and seven type I receptors (also known as Activin-receptor-Like-Kinases: ALKs 1-7). The TGF $\beta$  subfamily of ligands form unique receptor complexes by pairing specific type II receptors (TGF $\beta$ R-II or ActR-IIB) with specific type I (ALK4, ALK5 or ALK7) receptors. Similarly the BMP family of ligands construct receptor complexes by pairing specific type II receptors (BMPR-II or ActR-IIA/B or AMHR-II) with specific ALKs (ALK1, ALK2, ALK3 or ALK6) [1, 8]. Once activated ALKs 4, 5 and 7 primarily phosphorylate Smad2 and 3 while ALKs 1, 2, 3 and 6 phosphorylate Smads 1, 5, and 8 at the highly conserved C-terminal Ser-Xxx-Ser motif. This phosphorylation of dual residues, often referred to as tail-phosphorylation, triggers the binding of Smads to co-Smad4 and their translocation to the nucleus. In the nucleus Smad4 and tail-phosphorylated Smads form functional complexes with other cofactors and regulatory proteins and regulate the transcription of over 500 genes, which control context-specific cellular outcomes [1, 10].

Given the indispensable roles of ALKs in driving the TGF $\beta$  and BMP pathways, they have become attractive targets for the development of small molecule inhibitors to attenuate the cellular effects of TGF $\beta$  and BMP ligands. Among TGF $\beta$  ligands, TGF $\beta$  1-3, Activin and Nodal lead to the activation of ALK5, ALK4 and ALK7 respectively [1]. The kinase domains of ALKs 4, 5 and 7 are highly related to each other structurally. Similarly the kinase domains of BMP-activated ALKs (ALKs 1, 2, 3, and 6) display a high degree of similarity with one another, although among these ALK1 is more closely related to ALK2, and ALK3 is more closely related to ALK6 [11]. The expression of ALK1 is limited to certain cell types, primarily in endothelial cells, and has also been implicated in mediating TGF $\beta$ -induced phosphorylation of Smad1/5/8 in conjunction with ALK5 [12, 13]. Over the past few years several small molecule inhibitors of various ALKs have been developed. SB-431542 [11, 14, 15], SB-505124 [16], SB-525334 [17], LY-364947 [18, 19], A-83-01 [20], LY-2157299 [21], GW-6604 [22] and SD-208 [23] have all been reported as selective inhibitors of the TGF $\beta$ -activated ALKs (ALK4, 5 and 7). Similarly, more recently Compound C (but renamed Dorsomorphin to describe its effect in zebrafish embryos) [24] and its derivative LDN-

193189 [6, 25] have been reported as selective inhibitors of the BMP-activated ALKs (ALKs 2, 3 and 6). Although these compounds are described as “selective” inhibitors of specific ALKs, extensive specificity tests against other protein kinases have not been reported, except for Compound C [26]. Here we report the specificities and potencies of the most commonly used chemical inhibitors of the TGF $\beta$  pathway (SB-431542, SB-505124, LY-364947 and A-83-01) and BMP pathway (Dorsomorphin and LDN-193189) (Figure 1) by profiling these against a panel of up to 123 protein kinases covering a broad spectrum of the human kinome. Our data indicate that among the TGF $\beta$  pathway inhibitors, SB-505124 is a potent and selective inhibitor of TGF $\beta$ -activated ALKs. We also demonstrate that both of the BMP pathway inhibitors, Dorsomorphin and LDN-193189, can potently inhibit multiple protein kinases in addition to the BMP-activated ALKs. Our specificity data will be useful for researchers considering the use of these molecules as BMP and TGF $\beta$  pathway inhibitors.

## 2. Materials and Methods

### 2.1. Materials

SB-431542, SB-505124, LY-364947 and Dorsomorphin (Compound C) were purchased from Sigma. A-83-01 was purchased from Tocris Bioscience. LDN-193189 was purchased from Stemgent.  $^{32}$ P  $\gamma$ -ATP was from Perkin-Elmer. BMP-2 and TGF $\beta$ 1 were from R&D Biosystems. Meso-diaminopimelic acid (meso-DAP) was synthesized by Natalia Shpiro. DMSO and Tween-20 were from Sigma. Active GST-ALK2 and GST-ALK4 were purchased from Carina Biosciences. Antibodies recognising phospho-Smad1/5/8, phospho-Smad2, GAPDH, phospho-ERK1/2 and total ERK1/2 were from Cell Signalling.

### 2.2. General Methods

Tissue culture, immunoblotting, restriction enzyme digests, DNA ligations and other recombinant DNA procedures were performed using standard protocols. All DNA constructs used were verified by DNA sequencing, performed by DNA Sequencing & Services (MRCPPU, College of Life Sciences, University of Dundee, Scotland, [www.dnaseq.co.uk](http://www.dnaseq.co.uk)) using Applied Biosystems Big-Dye Ver 3.1 chemistry on an Applied Biosystems model 3730 automated capillary DNA sequencer.

### 2.3. Specificity Kinase panel

All protein kinases in the specificity panel were expressed, purified and assayed at The National Centre for Protein Kinase Profiling (<http://www.kinase-screen.mrc.ac.uk/>) as previously described [26]. Briefly, all assays (except ALK assays, which are described below) were carried out robotically at room temperature (21 °C) and were linear with respect to time and enzyme concentration under the conditions used. Assays were performed for 30 min using Multidrop Micro reagent dispensers (Thermo Electron Corporation, Waltham, MA, U.S.A.) in a 96-well format. The abbreviations for each of the kinases are defined in the legend to Figure 2. The concentration of magnesium acetate in the assays was 10 mM and [ $\gamma$ - $^{33}$ P]ATP (~800 cpm/pmol) was used at 5  $\mu$ M for ABL, Aurora A, CK2 $\alpha$ , CLK2, DAPK1, DYRK3, EF2K, EIF2AK3, ERK1, ERK8, GSK3 $\beta$ , HER4, HIPK2, IGF-1R, IKK $\beta$ , IRAK1, IRR, JAK2, MARK3, MKK1, MKK2, p38 $\alpha$  MAPK, p38 $\gamma$  MAPK, PAK2, PAK5, PIM3, PKB $\alpha$ , PKC $\zeta$ , PRAK, RIPK2, TAK1, TLK1 and ZAP70, 20  $\mu$ M for Aurora B, BRK, BRSK1, CAMKK $\beta$ , CDK2-Cyclin A, CHK1, CHK2, CK1 $\delta$ , CSK, EPH-B1, EPH-B2, EPH-B3, ERK2, FGF-R1, GSK, HIPK1, HIPK3, IR, IRAK4, JNK1, JNK2, JNK3, LKB1, MAPKAP-K2, MAPKAP-K3, MARK1, MARK2, MEKK1, MLK3, MNK1, MSK1, MST4, NEK2 $\alpha$ , OSR1, p38 $\delta$  MAPK, PAK4, PAK6, PDK1, PIM1, PIM2, PKA, PKC $\gamma$ , PKD1, PLK1, PRK2, ROCK2, RSK1, SGK1, SmMLCK, SYK, TAO1, TIE2, TrkA, TTK, VEG-FR and YES1 and 50  $\mu$ M for AMPK, ASK1, BRSK2, BTK, CAMK1, DYRK1A, DYRK2, EPH-A2, EPH-A4, EPH-B4, IKK $\epsilon$ , Lck, MARK4, MELK, MINK1, MKK6, MLK1, MNK2, MPSK1, MST2, NEK6, NUA1, p38 $\beta$  MAPK, PHK, PKB $\beta$ , PKC $\alpha$ , RSK2, S6K1, Src, SRPK1,

STK33 and TBK1 in order to be at or below the  $K_m$  for ATP for each enzyme. For kinase assays with CK1 isoforms, 300  $\mu$ M CK1-peptide KRRRALS\*VASLPGL (where S\* is phospho Serine) was used as the substrate.

#### 2.4. Cell Culture, Manipulation and Lysis:

Human keratinocyte (HaCaT) cells were cultured in 10-cm diameter dishes in Dulbecco's Modified Eagle's medium (DMEM) supplemented with 10% Foetal bovine serum, 1% penicillin/streptomycin mix and 2 mM L-Glutamine (D10F). RAW 264.7 cells (Mouse leukaemic monocyte macrophage cell line) were cultured as above except that foetal bovine serum was heat inactivated at 65 °C for 1h prior to use. Both cell lines were grown under a humidified atmosphere with 5 % CO<sub>2</sub> at a constant temperature of 37 °C. HaCaT cells were deprived of serum for 16h prior to treatment with ligands or inhibitors. Unless stated otherwise, cells were treated with the appropriate small molecule inhibitors or solvent control 2h prior to treating cells with BMP-2 (25 ng/ml final), TGF $\beta$  (50 pM final) or Meso-DAP (15  $\mu$ M) for 1h. Cells were then washed once with ice-cold PBS and lysed in 0.5 ml ice-cold complete lysis buffer (50 mM Tris-HCl pH7.5, 1 mM EGTA, 1 mM EDTA, 1 % Triton X-100, 1 mM sodium orthovanadate, 50 mM sodium fluoride, 5mM sodium pyrophosphate, 0.27 M sucrose, 5 mM  $\beta$ -glycerophosphate, 0.1 % (v/v) 2-mercaptoethanol, 1 tablet per 25 ml of complete protease inhibitor cocktail). The extracts were spun down at 16,000 g at 4 °C for 10 minutes prior to snap-freezing in liquid nitrogen and storing at -80 °C if not processed immediately.

#### 2.5. SDS-PAGE and Western Blotting:

Cell extracts (20  $\mu$ g) were heated at 95 °C for 5 min in 1XSDS sample buffer (62.5 mM Tris-HCl pH 6.8, 10% (v/v) Glycerol, 2% (w/v) SDS, 0.02% (w/v) bromophenol blue, and 1% (v/v)  $\beta$ -mercaptoethanol), resolved on a 10% polyacrylamide gel by electrophoresis and transferred to nitrocellulose membranes. Membranes were blocked in TBS-T buffer [50 mM Tris-HCl pH 7.5, 0.15 M NaCl and 0.1% (w/v) Tween-20] containing 10% (w/v) non-fat milk. The membranes were then incubated with the indicated antibodies, diluted in TBS-T containing 10% (w/v) milk for 16 h at 4 °C. The membranes were washed 2x10 min in TBS-T buffer, probed with the secondary antibody (either HRP-conjugated or IRDye-800 or -680 conjugated and diluted 1:5000 in TBS-T/5% milk) for 1h at room temperature, and washed 3x10 min in TBS-T buffer. Detection was performed by using enhanced chemiluminescence reagent for HRP-conjugated secondary antibodies and by using the Odyssey Imaging System (LI-COR Biosciences) for IRDye-800 or -680 conjugated antibodies. For IC<sub>50</sub> determinations, the intensities of the bands corresponding to appropriate phosphorylated Smads and corresponding total Smads were quantified using the Odyssey Imaging System software.

2.6. *ALK2, ALK3, ALK4 and ALK5 Kinase assays:* N-terminal GST-tagged constitutively active mutant of ALK3 (Q233D, 187-532), wild type GST-ALK5 (200-503) and wild type GST-BMPRII (174-1038) were cloned into pFastBac baculovirus vectors (Invitrogen) and expressed in Sf9 insect cells. For kinase assays, 20  $\mu$ l reactions were setup consisting of 150 ng of kinase (GST-ALK2, 3, 4 or 5) and 2  $\mu$ g substrate protein (GST-Smad1 or 2) in a buffer containing 50 mM TrisHCl pH 7.5, 0.1% 2-mercaptoethanol, 0.1 mM EGTA, 10 mM MgCl<sub>2</sub>, 0.5  $\mu$ M Microcystin-LR, 0.1 mM  $\gamma$ -<sup>32</sup>P-ATP (500 cpm/pmole) and 5% DMSO or DMSO containing the appropriate concentrations of the small molecule inhibitors. For ALK2 and ALK3 assays, GST-Smad1 was used as a substrate. For ALK4 and ALK5 assays, GST-Smad2 was used as a substrate. ALK3 assays also contained 150 ng of GST-BMPRII. The assays were performed at 30 °C for 30 minutes and stopped by adding 1X SDS sample buffer and heating at 95 °C for 5 minutes. The samples were resolved by SDS-PAGE, the gels stained with Coomassie Blue and dried. The radioactivity was analysed by autoradiography. For IC<sub>50</sub> and percentage kinase activity remaining determinations, the stained bands representing protein substrates were excised and the radioactivity measured.

### 3. Results

#### 3.1. Specificities of Inhibitors of the TGF $\beta$ pathway

Although multiple small molecule inhibitors have been reported as specific inhibitors of TGF $\beta$  pathway, SB-431542, SB-505124, LY-364947 and A-83-01 have been the most widely exploited in studies investigating the physiological roles of TGF $\beta$  ligands, together accounting for over 300 publications. In this study we have profiled the specificities of these molecules against a panel of up to 123 protein kinases *in vitro* (Figure 2A-D; Figure 4). Structurally SB-431542 is similar to SB-505124 and LY-364947 is similar to A-83-01 (Figure 1).

#### 3.2. Specificities and Potencies of SB-431542 and SB-505124

SB-431542 and SB-505124 were both developed as ALK5 inhibitors from triarylimidazole templates [11, 14-16]. Both compounds are ATP-competitive, reversible inhibitors of ALK5 and can also inhibit ALKs 4 and 7 [11, 14-16]. SB-431542 was the first small molecule inhibitor of ALKs 4, 5 and 7 to be reported and has been the most widely used inhibitor of the TGF $\beta$  pathway resulting in over 200 research reports [11, 14, 15]. We tested the ability of both SB-431542 and SB-505124 to inhibit the activity of a panel of over 105 protein kinases at two different concentrations (Figure 2A&B). At 1  $\mu$ M, besides ALK5, SB-431542 inhibited RIPK2 and CK1 $\delta$  activities by 77% and 70% respectively, while p38 $\alpha$  MAPK was inhibited by 30% (Figure 2A). At 0.1  $\mu$ M, SB-431542 inhibited RIPK2 and CK1 $\delta$  by 33% and 29% respectively (Figure 2A). Similarly 1  $\mu$ M SB-505124 inhibited RIPK2 by about 72% and p38 $\alpha$  MAPK by 49% but did not inhibit CK1 $\delta$  (Figure 2B). At 0.1  $\mu$ M, SB-505124 inhibited RIPK2 by 18% but CK1 $\delta$  and p38 $\alpha$  MAPK were not inhibited (Figure 2B). At both concentrations, SB-431542 and SB-505124 inhibited ALK5 activity *in vitro* but did not inhibit ALK3 (Figure 2A&B, Figure 4A&B). At both concentrations the activities of all other kinases in the panel were not significantly inhibited by either of these compounds (Figure 2A&B).

SB-431542 inhibits the phosphorylation of Smad3 by ALK5 and ALK4 *in vitro* with an IC<sub>50</sub> of 0.094  $\mu$ M and 0.14  $\mu$ M respectively [11] (Table 1A). In contrast, SB-505124 inhibits the phosphorylation of Smad3 by ALK5 and ALK4 with an IC<sub>50</sub> of 0.047  $\mu$ M and 0.129  $\mu$ M respectively [16] (Table 1A). We determined that SB-431542 and SB-505124 inhibit RIPK2 with IC<sub>50</sub> of 0.41  $\mu$ M (4-fold lower potency than that seen for ALK5) and 0.35  $\mu$ M (7-fold) respectively (Table 2). Because SB-431542 also potently inhibited CK1 $\delta$  at 1  $\mu$ M (Figure 2A), we tested the ability of both SB-431542 and SB-505124 to inhibit CK1 isoforms *in vitro*. SB-431542 potently inhibited CK1 $\alpha$ , CK1 $\delta$  and CK1 $\epsilon$  isoforms with IC<sub>50</sub> of 1.34  $\mu$ M, 0.92  $\mu$ M and 0.38  $\mu$ M respectively but did not inhibit CK1 $\gamma$  (Table 2). SB-505124 inhibited CK1 $\alpha$ , CK1 $\delta$  and CK1 $\epsilon$  isoforms with IC<sub>50</sub> of 19.44  $\mu$ M, 3.38  $\mu$ M and 1.60  $\mu$ M respectively but did not inhibit CK1 $\gamma$  (Table 2). Both SB-431542 and SB-505124 also inhibit p38 $\alpha$  MAPK at high concentrations with IC<sub>50</sub> values reported to be >10  $\mu$ M [11, 16].

SB-505124 is reported to be a more potent inhibitor of the TGF $\beta$  pathway in cells than SB-431542 [16]. In multiple cell lines the TGF $\beta$  induced phosphorylation of Smad2 was inhibited by SB-505124 and SB-431542 with IC<sub>50</sub> values of ~0.25  $\mu$ M and 0.5-1  $\mu$ M respectively [11, 16] (Table 1A). Similarly SB-505124 inhibited the ability of constitutively active ALK5 to induce the expression of CAGA-luciferase reporter activity more potently than SB-431542 [16] (Table 1A). Both inhibitors were also shown to inhibit the phosphorylation of Smad2 and expression of CAGA-Luciferase reporter activities driven by constitutively active ALK4 and ALK7 [11, 16].

#### 3.3. Specificities and Potencies of LY-364947 and A-83-01

LY-364947, a pyrazole-based small molecule, was developed as an inhibitor of ALK5 and is an ATP-competitive, cell permeable inhibitor [18, 19]. *In vitro*, it inhibits ALK5 with an IC<sub>50</sub>

of 0.058  $\mu\text{M}$ , potency comparable to that of SB-505124 [19](Table 1A). Furthermore it inhibits TGF $\beta$ -induced phosphorylation of Smad2 in cells with a similar potency as SB-505124 (Table 1A). When profiled against 123 protein kinases (Figure 2C), LY-364947 at 1  $\mu\text{M}$  inhibited ALK5, RIPK2, VEGF-R, CK1 $\delta$  and MINK1 activity by more than 50% and at 10  $\mu\text{M}$ , in addition to these kinases, it inhibited p38 $\alpha$  MAPK, PKD1, GCK, BRK, Lck, TAK1, YES1, FGF-R1 and p38 $\beta$  MAPK by more than 50%. LY-364947 inhibited RIPK2 and CK1 $\delta$  with IC<sub>50</sub> of 0.11  $\mu\text{M}$  and 0.22  $\mu\text{M}$  respectively (Table 2). Similarly LY-364947 would be predicted to inhibit VEGF-R and MINK1 with similar IC<sub>50</sub> values (Figure 2C). LY-364947 inhibited CK1 $\alpha$ , CK1 $\epsilon$  and CK1 $\gamma$  isoforms with IC<sub>50</sub> of 2.27  $\mu\text{M}$ , 1.34  $\mu\text{M}$  and 44  $\mu\text{M}$  respectively (Table 2).

A-83-01, structurally related to LY-364947 (Figure 1), was developed as an inhibitor of the TGF $\beta$  pathway using a cell-based CAGA-Luciferase reporter assay driven by constitutively active ALKs4, 5 and 7 [20]. A-83-01 inhibited TGF $\beta$ -induced CAGA-Luciferase reporter activity in Mv1Lu lung epithelial cells with an IC<sub>50</sub> of 0.03  $\mu\text{M}$  [20](Table 1A), more potently than SB-431542 (IC<sub>50</sub>, 0.25  $\mu\text{M}$ ) and SB-505124 (IC<sub>50</sub>, 0.1  $\mu\text{M}$ ). However detailed kinetic analysis of the ability of A-83-01 to inhibit different ALKs *in vitro* has not been reported [20]. Nonetheless we tested the ability of A-83-01 to inhibit a panel of 107 kinases at 1  $\mu\text{M}$  and 0.1  $\mu\text{M}$  (Figure 2D). We demonstrate that at 1  $\mu\text{M}$ , A-83-01 inhibited ALK5, VEG-FR, RIPK2, MINK1, p38 $\alpha$  MAPK, PKD1 and FGF-R1 by more than 50% (Figure 2D). At 0.1  $\mu\text{M}$ , ALK5, VEG-FR, RIPK2 were inhibited by more than 50% while MINK1, p38 $\alpha$  MAPK and FGF-R1 were inhibited by more than 30% (Figure 2D). A-83-01 potently inhibited RIPK2 with an IC<sub>50</sub> of 0.1  $\mu\text{M}$  (Table 1B) and would be predicted to inhibit VEGF-R with similar potency (Figure 2D). A-83-01 inhibited CK1 $\alpha$ , CK1 $\delta$ , CK1 $\epsilon$  and CK1 $\gamma$  isoforms with IC<sub>50</sub> of 15.66  $\mu\text{M}$ , 3.42  $\mu\text{M}$ , 4.59  $\mu\text{M}$  and 29  $\mu\text{M}$  respectively (Table 2).

### 3.4. Specificities of Inhibitors of the BMP pathway

Recently, Dorsomorphin (Compound C) and LDN-193189, a Dorsomorphin derivative, were reported as selective and potent inhibitors of the BMP pathway [6, 24, 25]. Subsequently these compounds have been widely used in cell-based assays and whole organisms to study the physiological roles of the BMP pathway. In this study we have profiled the specificities of these molecules against a panel of up to 121 protein kinases *in vitro* (Figure 3A&B).

### 3.5 Specificity of Dorsomorphin (Compound C) as a BMP pathway inhibitor

In vertebrates BMP signalling plays a crucial role in defining dorso-ventral (DV) axis, where inhibition of BMP pathway results in dorsalised axis patterning [27]. A high throughput small molecule screen in zebrafish embryos identified Compound C (this was renamed Dorsomorphin) as an inhibitor of the BMP pathway as it resulted in dorsalised axis patterning of zebrafish embryos [24]. Subsequently Dorsomorphin was reported as a selective small molecule inhibitor of BMP pathway and was shown to inhibit BMP-activated ALKs 2, 3 and 6 [24]. Previously Compound C has been described, and extensively used, as a selective inhibitor of AMPK [28]. However, a study looking at the specificity of Compound C profiled against a panel of 70 kinases found that it inhibited a number of kinases, including ERK8, MNK1, PHK, MELK, DYRK isoforms, HIPK2, Src and Lck, with similar or greater potency than AMPK [26]. This information on the specificity profile of Compound C [26] has been overlooked by all the reports describing or employing Dorsomorphin as a specific inhibitor of the BMP pathway [24, 29-31].

In this study we extended the specificity and the potency tests on Dorsomorphin at three different concentrations against a panel of 119 protein kinases (Figure 3A). At 10  $\mu\text{M}$ , Dorsomorphin inhibited the activities of 64 out of the 119 kinases by >50%. At 1  $\mu\text{M}$ , Dorsomorphin inhibited the activities of 34 out of 119 kinases more potently than it inhibited AMPK and by >50% (Figure 3A). Even at 0.1  $\mu\text{M}$ , VEGF-R, ERK8, GCK, CLK2, DYRK1A, PHK, ABL, NUA1, PRK2 and YES1 were inhibited by >50% implying that Dorsomorphin inhibits these kinases with IC<sub>50</sub> values lower than 0.1  $\mu\text{M}$  (Figure 3A). Dorsomorphin



inhibited ALK3 *in vitro* with an IC<sub>50</sub> of ~1 μM while it did not inhibit ALK5 *in vitro* (Figure 4C). Furthermore, Dorsomorphin was reported to inhibit the BMP Responsive Element (BRE)-Luciferase reporter activity driven by constitutively active ALK2, ALK3 and ALK6 with IC<sub>50</sub> values of 0.2 μM, 0.5 μM and 5-10 μM respectively [24] (Table 1B). Clearly Dorsomorphin is not a selective inhibitor of the BMP pathway and is therefore not a good candidate for selective inhibition of BMP-activated ALKs. In fact it was recently reported that in zebrafish, Dorsomorphin, when used at concentrations sufficient to inhibit the BMP pathway, strongly inhibited intersegmental vessel formation by inhibiting VEGF-R2 [32], demonstrating the potential off-target effects of using a non-selective inhibitor.

### 3.6. Specificity of LDN-193189 as a BMP pathway inhibitor

Using Dorsomorphin as a template, LDN-193189 was developed as a compound with improved potency as a BMP pathway inhibitor [25]. In rat pulmonary artery smooth muscle cells (rPASC), it was shown that LDN-193189 inhibited BMP4-induced phosphorylation of Smad1/5/8 with an IC<sub>50</sub> of 0.005 μM, an improvement of 94-fold over Dorsomorphin, which it inhibits with an IC<sub>50</sub> of 0.47 μM [25]. In human keratinocyte HaCaT cells, we observed that LDN-193189 inhibits BMP2-induced phosphorylation of Smad1/5/8 with an IC<sub>50</sub> of ~0.005 μM (Figure 5A). In contrast the TGFβ-induced phosphorylation of Smad2 was only slightly affected at >3 μM but TGFβ-induced phosphorylation of Smad1/5/8 was inhibited robustly at 0.3 μM LDN-193189 (Figure 5B). We next assessed the ability of LDN-193189 to inhibit various ALKs *in vitro* using 100 μM ATP in the assays (Figure 6). LDN-193189 inhibited the ability of ALK2 to phosphorylate GST-Smad1 *in vitro* with an IC<sub>50</sub> of 45 nM, while its autophosphorylation was inhibited with an IC<sub>50</sub> of 30 nM (Figure 6A). LDN-193189 inhibited the ability of ALK3 to phosphorylate Smad1 *in vitro* with an IC<sub>50</sub> of 100 nM, although even at 3 μM, ALK3 was not completely inhibited (Figure 6B). It is also noteworthy that autophosphorylation of BMPRII, which was also present in the assay and is required in order to activate ALK3 (data not shown), was not inhibited by LDN-193189, implying that BMPRII is not inhibited by LDN-193189 (Figure 6B). LDN-193189 inhibited ALK4 and ALK5 with much higher IC<sub>50</sub> values of 0.3 μM and 0.5 μM respectively (Figure 6C&D). LDN-193189 has been reported to inhibit the BRE-Luciferase reporter activity driven by constitutively active ALK2 and ALK3 with IC<sub>50</sub> of 0.005 μM and 0.03 μM respectively [6] (Table 1B). Overall it is evident that LDN-193189 is a very potent inhibitor of the BMP pathway and that it can inhibit BMP-activated ALK2 and ALK3 *in vitro*. However, despite studies describing LDN-193189 as a selective and potent inhibitor of the BMP pathway, its specificity and potency has not been tested against an extensive array of protein kinases.

We profiled the specificity and potency of LDN-193189 at three different concentrations against a panel of 121 protein kinases covering a broad spectrum of the human kinome (Figure 3B). We noted a very similar specificity and potency profile for LDN-193189 and Dorsomorphin (Figure 3A&B). Like Dorsomorphin, we found that at 10 μM, LDN-193189 inhibited 44 out of the 121 kinases by >50%, majority of them very potently (Figure 3B). At 1 μM, LDN-193189 inhibited 24 out of the 121 protein kinases by >50% and of these RIPK2, FGF-R1, NUA1, CAMKKβ, MINK1, GSK, VEGFR, BRK, YES1 and CLK2 were inhibited very potently. Even at 0.1 μM, LDN-193189 inhibited RIPK2, FGF-R1, NUA1, CAMKKβ, MINK1, GSK, VEGFR and BRK by >50%, implying that these kinases are inhibited by LDN-193189 with IC<sub>50</sub> values lower than 0.1 μM (Figure 3B). Indeed LDN-193189 inhibits RIPK2 and GSK with IC<sub>50</sub> values of 0.025 μM and 0.08 μM respectively, values similar to those seen against ALK2 and ALK3 *in vitro* respectively (Table 2; Figure 6A&B).

### 3.7. Inhibition of RIPK2 by LDN-193189 in RAW macrophage cells

We noted from above that RIPK2, a member of the receptor interacting protein (RIP) family of protein kinases, was inhibited potently *in vitro* by LDN-193189 (Table 2). RIPK2 is implicated in NOD1 and NOD2 signalling and results in the activation of MAP Kinases, NFκB and inflammatory mediators in response to NOD1/2 agonists. In cells, NOD1 and

NOD2 sense peptidoglycan-related molecules from intracellular bacteria that have evaded recognition by Toll-like receptors at the cell-surface [33]. In RAW macrophages cells, the peptidoglycan-related agonist D-glutamyl-meso-diaminopimelic acid (meso-DAP) induced the phosphorylation of ERK1/2 (Figure 7). This induction was inhibited when cells were treated with 0.1  $\mu$ M or 10  $\mu$ M LDN-193189 (Figure 7), concentrations that are sufficient to inhibit BMP signalling (Figure 5A).

## 4. Discussion

In this report we have examined the specificities of the most commonly used small molecule inhibitors of the TGF $\beta$  and BMP pathways against a panel of up to 123 protein kinases. The results highlight the potential off-target effects of these small molecule inhibitors when using them to assess the physiological roles of TGF $\beta$  and BMP pathways. Furthermore, because the specificity was profiled against only 23% of the human protein kinases, these inhibitors are likely to inhibit other kinases that have not yet been studied. While caution is recommended in interpreting any impact on the TGF $\beta$  and BMP pathways resulting from the use of these inhibitors, the specificity profile provided herein should provide useful information for researchers when deciding which inhibitor to use. We also emphasize the fact that the specificity profiles presented in this report were obtained using *in vitro* kinase assays. We recommend using the minimum effective-concentrations against intended targets when using any chemical inhibitors to inhibit the BMP/TGF $\beta$  pathways in cell and animal based assays and to test thoroughly whether at these concentrations the molecules also inhibit other kinases that are inhibited potently *in vitro*.

### 4.1. Inhibitors of the TGF $\beta$ pathway:

Active TGF $\beta$  signalling has been implicated in the development of fibrotic sclerosis of multiple organs including heart, kidney, lungs, liver and skin [2-4]. TGF $\beta$  signalling is also associated with promotion of cancer progression and metastasis [5, 34]. As a result, TGF $\beta$  - activated ALKs, in particular ALK5, have been targeted for the development of small molecule inhibitors by major pharmaceutical industries [14, 18, 35]. Many ALK5 inhibitors, which also potently inhibit ALK4 and ALK7, have entered pre-clinical trials to treat fibrosis and advanced metastatic cancers and have met with mixed results [35]. The specificity of chemical inhibitors is particularly important when using them in whole organisms, as consequences of off-target effects could lead to undesirable side effects.

Based on specificity and potency of the four inhibitors of the TGF $\beta$  pathway, we recommend the use of SB-505124, at or below 1 $\mu$ M, as an inhibitor of ALKs 4, 5 and 7 in cell based assays. While both SB-431542 and SB-505124 are relatively selective inhibitors of ALKs 4, 5 and 7, SB-505124 is a more potent inhibitor of ALK4, 5 and 7 and inhibits CK1 isoforms less potently than SB-431542 (Table 1A & 2). Furthermore, in cell-based assays, SB-505124 was reported to be less cytotoxic than SB-431542 [16]. Both inhibit RIPK2 with similar IC<sub>50</sub> values (Table 2) and we recommend that RIPK2 inhibition be assessed at concentrations of SB-505124 used to inhibit TGF $\beta$  signalling. SB-525334, which is structurally very closely related to SB-505124 (Figure 1), has been reported to be around 3-fold more potent inhibitor of ALK5 and ALK4 compared to SB-505124 [36], however it has not been used as extensively as other ALK5 inhibitors. At concentrations sufficient to inhibit ALK5, both LY-364947 and A-83-01 inhibited RIPK2, MINK1 and VEGF-R potently. LY-364947 also inhibited CK1 isoforms potently while A-83-01 inhibited p38 $\alpha$  MAPK, PKD1 and FGF-R1 potently (Figure 2C&D). When using LY-364947 and A-83-01 as TGF $\beta$  pathway inhibitors, these potential off-target effects have to be considered. One of the impediments to using small molecule inhibitors of TGF $\beta$  pathway is that they inhibit ALKs 4, 5 and 7 and show no significant selectivity between these ALKs. Knockout models of ALK4, ALK5 or ALK7 display unique phenotypes [37, 38] suggesting unique cellular or contextual roles for these ALKs. Development of ALK-specific inhibitors will be essential to probe the roles of

individual ALKs in cells as well as target selective ALKs that may be responsible for driving a particular disease states.

#### 4.2. Inhibitors of the BMP pathway:

While strong pharmaceutical efforts to develop small molecule inhibitors against the TGF $\beta$  pathway have led to multiple ALK4, ALK5 and ALK7 inhibitors, the development of small molecule inhibitors of the BMP pathway has lagged behind. BMP signalling plays critical roles during embryogenesis, in controlling the fate of various progenitor cell populations, including embryonic stem cells and hematopoietic stem cells, and in most differentiated and specialized cells and in skeletogenesis [39-42]. Selective small molecule inhibitors of the BMP pathway are desirable in dissecting the physiological roles of BMP signalling in different cellular contexts. ALKs 2, 3 and 6 mediate BMP signals in most tissues, while ALK1, expressed mainly in endothelial cells, signals through both BMP and TGF $\beta$  ligands [12, 13]. Sustained BMP signalling driven by a constitutively active mutants of ALK6 have been implicated in heterotopic ossification [6]. Similarly overexpression of certain BMP ligands and activation of downstream signalling has been reported in some cancers [43]. Selective small molecule inhibitors of BMP activated ALKs could be therapeutically beneficial against these diseases. Dorsomorphin and LDN-193189, the only two small molecule inhibitors of the BMP pathway are rather non-specific as they inhibit a number of other protein kinases potently. Some of the off-target effects of using Dorsomorphin at concentrations sufficient to inhibit BMP signalling have been demonstrated and others likely exist [26, 32]. For these reasons use of Dorsomorphin to inhibit BMP pathway is not recommended.

LDN-193189 on the other hand is a very potent inhibitor of BMP signalling, inhibiting BMP-induced phosphorylation of Smad1 in cells with an IC<sub>50</sub> of 5 nM [6, 25] (Figure 5A). Because ALK2 and ALK3 were inhibited *in vitro* by LDN-193189 with IC<sub>50</sub> of 30-45 nM and 100 nM respectively, it was rather surprising that the BMP pathway in cells was inhibited with a substantially lower IC<sub>50</sub>. This could mean that LDN-193189 binds very efficiently, possibly allosterically, to the BMP-activated ALKs in cells. However crystal structures of LDN-193189 in complex with the kinase domain of ALK1 shows LDN-193189 binding to the ATP-binding pocket of ALK1 kinase domain (Link: [http://www.thesgc.org/structures/structure\\_description/3MY0/](http://www.thesgc.org/structures/structure_description/3MY0/)). Alternatively the binding of LDN-193189 to BMP-activated ALKs could affect the formation of BMP-induced complexes between these ALKs and the upstream type II receptors, which would thus inhibit the activation of type I receptors. Furthermore like Dorsomorphin, LDN-193189 inhibited a number of other kinases very potently. While LDN-193189 displayed improved potency against BMP-activated ALKs over Dorsomorphin, the potency with which both of these compounds inhibit many other kinases did not change significantly (Figure 3). Therefore it is possible that the effects on LDN-193189 to inhibit BMP signalling so potently could be only partly dependent on its effects on ALKs and partly on its effects on other protein kinases, which may impact on the activation or activity of ALKs or the access of ALKs to their substrates. When using LDN-193189 as an inhibitor of the BMP pathway in cells or whole organisms, the consequences of its ability to inhibit other kinases, notably RIPK2, FGF-R1, NUAK1, CAMKK $\beta$ , MINK1, GCK, VEG-FR and BRK should be considered. Nonetheless due to its potency as a BMP pathway inhibitor, LDN-193189 provides a very good platform to design derivatives that could enhance its selectivity for BMP-activated ALKs.

#### Conclusions:

- Based on *in vitro* specificity tests against a substantial panel of human kinases, routinely used TGF $\beta$  pathway inhibitors (SB-431542, SB-505124, LY-364947 and A-83-01) are relatively more selective than the BMP pathway inhibitors (Dorsomorphin and LDN-193189).

- Of the TGF $\beta$  pathway inhibitors, we recommend SB-505124, to be used at or below 1 $\mu$ M for cell-based assays, as the most suitable molecule for use as an inhibitor of ALKs 4, 5 & 7 and the TGF $\beta$  pathway.
- Dorsomorphin, also known as Compound C, is a non-specific inhibitor of BMP-activated ALKs and potentially inhibits many more kinases *in vitro*. Therefore we do not recommend the use of Dorsomorphin as a selective BMP pathway inhibitor.
- LDN-193189, while being a potent inhibitor of the BMP pathway, also potentially inhibits multiple kinases *in vitro*. At concentrations sufficient to inhibit the BMP pathway, LDN-193189 inhibits RIPK2-mediated phosphorylation of ERK1/2 in cells. Therefore its use as a selective inhibitor of the BMP pathway has to be considered with caution.

### Acknowledgements:

We thank Thomas Macartney for the cloning of ALK3, ALK5, BMPR2 and Smad constructs. We thank the staff at the National Centre for Protein Kinase Profiling ([www.kinase-screen.mrc.ac.uk](http://www.kinase-screen.mrc.ac.uk)) for undertaking the kinase specificity screening, the Sequencing Service (School of Life Sciences, University of Dundee, Scotland) for DNA sequencing and the protein production teams [Division of Signal Transduction Therapy (DSTT), University of Dundee] coordinated by Hilary McLauchlan and James Hastie for expression and purification of proteins. We thank Mazin Al-Salihi and David Bruce for helpful discussions. We thank the Medical Research Council, and the pharmaceutical companies supporting the Division of Signal Transduction Therapy Unit (AstraZeneca, Boehringer-Ingelheim, GlaxoSmithKline, Merck-Serono and Pfizer) for financial support.

### References

- [1] Shi Y, Massague J, Cell. 2003;113:685-700.
- [2] Border WA, Noble NA, N Engl J Med. 1994;331:1286-1292.
- [3] Bottinger EP, Bitzer M, J Am Soc Nephrol. 2002;13:2600-2610.
- [4] Gu L, Zhu YJ, Yang X, Guo ZJ, Xu WB, Tian XL, Acta Pharmacol Sin. 2007;28:382-391.
- [5] Massague J, Cell. 2008;134:215-230.
- [6] Yu PB, Deng DY, Lai CS, Hong CC, Cuny GD, Boussein ML, Hong DW, McManus PM, Katagiri T, Sachidanandan C, Kamiya N, Fukuda T, Mishina Y, Peterson RT, Bloch KD, Nat Med. 2008;14:1363-1369.
- [7] Bierie B, Moses HL, Cytokine Growth Factor Rev. 21:49-59.
- [8] Heldin CH, Miyazono K, ten Dijke P, Nature. 1997;390:465-471.
- [9] Wrana JL, Attisano L, Wieser R, Ventura F, Massague J, Nature. 1994;370:341-347.
- [10] Massague J, Gomis RR, FEBS Lett. 2006;580:2811-2820.
- [11] Inman GJ, Nicolas FJ, Callahan JF, Harling JD, Gaster LM, Reith AD, Laping NJ, Hill CS, Mol Pharmacol. 2002;62:65-74.
- [12] Oh SP, Seki T, Goss KA, Imamura T, Yi Y, Donahoe PK, Li L, Miyazono K, ten Dijke P, Kim S, Li E, Proc Natl Acad Sci U S A. 2000;97:2626-2631.
- [13] Wrighton KH, Lin X, Yu PB, Feng XH, J Biol Chem. 2009;284:9755-9763.
- [14] Callahan JF, Burgess JL, Fornwald JA, Gaster LM, Harling JD, Harrington FP, Heer J, Kwon C, Lehr R, Mathur A, Olson BA, Weinstock J, Laping NJ, J Med Chem. 2002;45:999-1001.

- [15] Laping NJ, Grygielko E, Mathur A, Butter S, Bomberger J, Tweed C, Martin W, Fornwald J, Lehr R, Harling J, Gaster L, Callahan JF, Olson BA, *Mol Pharmacol.* 2002;62:58-64.
- [16] DaCosta Byfield S, Major C, Laping NJ, Roberts AB, *Mol Pharmacol.* 2004;65:744-752.
- [17] Laping NJ, Everitt JI, Frazier KS, Burgert M, Portis MJ, Cadacio C, Gold LI, Walker CL, *Clin Cancer Res.* 2007;13:3087-3099.
- [18] Sawyer JS, Anderson BD, Beight DW, Campbell RM, Jones ML, Herron DK, Lampe JW, McCowan JR, McMillen WT, Mort N, Parsons S, Smith EC, Vieth M, Weir LC, Yan L, Zhang F, Yingling JM, *J Med Chem.* 2003;46:3953-3956.
- [19] Peng SB, Yan L, Xia X, Watkins SA, Brooks HB, Beight D, Herron DK, Jones ML, Lampe JW, McMillen WT, Mort N, Sawyer JS, Yingling JM, *Biochemistry.* 2005;44:2293-2304.
- [20] Tojo M, Hamashima Y, Hanyu A, Kajimoto T, Saitoh M, Miyazono K, Node M, Imamura T, *Cancer Sci.* 2005;96:791-800.
- [21] Bueno L, de Alwis DP, Pitou C, Yingling J, Lahn M, Glatt S, Troconiz IF, *Eur J Cancer.* 2008;44:142-150.
- [22] de Gouville AC, Boullay V, Krysa G, Pilot J, Brusq JM, Lorient F, Gauthier JM, Papworth SA, Laroze A, Gellibert F, Huet S, *Br J Pharmacol.* 2005;145:166-177.
- [23] Uhl M, Aulwurm S, Wischhusen J, Weiler M, Ma JY, Almirez R, Mangadu R, Liu YW, Platten M, Herrlinger U, Murphy A, Wong DH, Wick W, Higgins LS, Weller M, *Cancer Res.* 2004;64:7954-7961.
- [24] Yu PB, Hong CC, Sachidanandan C, Babitt JL, Deng DY, Hoyng SA, Lin HY, Bloch KD, Peterson RT, *Nat Chem Biol.* 2008;4:33-41.
- [25] Cuny GD, Yu PB, Laha JK, Xing X, Liu JF, Lai CS, Deng DY, Sachidanandan C, Bloch KD, Peterson RT, *Bioorg Med Chem Lett.* 2008;18:4388-4392.
- [26] Bain J, Plater L, Elliott M, Shpiro N, Hastie CJ, McLauchlan H, Klevernic I, Arthur JS, Alessi DR, Cohen P, *Biochem J.* 2007;408:297-315.
- [27] Nguyen VH, Schmid B, Trout J, Connors SA, Ekker M, Mullins MC, *Dev Biol.* 1998;199:93-110.
- [28] Zhou G, Myers R, Li Y, Chen Y, Shen X, Fenyk-Melody J, Wu M, Ventre J, Doeber T, Fujii N, Musi N, Hirshman MF, Goodyear LJ, Moller DE, *J Clin Invest.* 2001;108:1167-1174.
- [29] Hong CC, Yu PB, *Cytokine Growth Factor Rev.* 2009;20:409-418.
- [30] Boergemann JH, Kopf J, Yu PB, Knaus P, *Int J Biochem Cell Biol.* 42:1802-1807.
- [31] Hao J, Daleo MA, Murphy CK, Yu PB, Ho JN, Hu J, Peterson RT, Hatzopoulos AK, Hong CC, *PLoS One.* 2008;3:e2904.
- [32] Cannon JE, Upton PD, Smith JC, Morrell NW, *Br J Pharmacol.* 161:140-149.
- [33] Inohara, Chamailard, McDonald C, Nunez G, *Annu Rev Biochem.* 2005;74:355-383.
- [34] Padua D, Massague J, *Cell Res.* 2009;19:89-102.
- [35] Seoane J, *Clin Transl Oncol.* 2008;10:14-19.
- [36] Grygielko ET, Martin WM, Tweed C, Thornton P, Harling J, Brooks DP, Laping NJ, *J Pharmacol Exp Ther.* 2005;313:943-951.
- [37] Itoh F, Itoh S, Carvalho RL, Adachi T, Ema M, Goumans MJ, Larsson J, Karlsson S, Takahashi S, Mummery CL, Dijke PT, Kato M, *Lab Invest.* 2009;89:800-810.

- [38] Jornvall H, Reissmann E, Andersson O, Mehrkash M, Ibanez CF, Mol Cell Biol. 2004;24:9383-9389.
- [39] Chen D, Zhao M, Mundy GR, Growth Factors. 2004;22:233-241.
- [40] De Robertis EM, Kuroda H, Annu Rev Cell Dev Biol. 2004;20:285-308.
- [41] Li X, Cao X, Ann N Y Acad Sci. 2006;1068:26-40.
- [42] Varga AC, Wrana JL, Oncogene. 2005;24:5713-5721.
- [43] Blanco Calvo M, Bolos Fernandez V, Medina Villaamil V, Aparicio Gallego G, Diaz Prado S, Grande Pulido E, Clin Transl Oncol. 2009;11:126-137.

## FIGURE LEGENDS

**Figure 1.** Chemical structures of the small molecule inhibitors of the TGF $\beta$  (A-E) and BMP pathways (F&G).

**Figure 2.** The specificities of TGF $\beta$  pathway inhibitors SB-431542 (A), SB-505124 (B), LY-364947 (C) & A-83-01 (D) against a panel of up to 123 protein kinases. The results are presented as bars indicating percentage activity remaining for each kinase (averages of two duplicate determinations) in the presence of the indicated concentration of the inhibitor compared with a control lacking the inhibitor  $\pm$  standard deviation. Further details of the assays are given in the Methods section. The results are ranked according to the percentage activity remaining when the assays were performed in the presence of the indicated inhibitors at 1  $\mu$ M. Protein kinases referred to in the Results section are indicated by red arrows.

Abbreviations for the protein kinases not described in the text are as follows: ABL, Abelson murine leukemia viral oncogene homolog; AMPK, AMP-activated protein kinase; ASK, Apoptosis signal regulating kinase; BRK, Breast tumour kinase; BRSK, brain-specific kinase; BTK, Bruton agammaglobulinemia tyrosine kinase; CaMK, calmodulin-dependent kinase; CaMKK, CaMK kinase; CDK, cyclin dependent kinase; CHK, checkpoint kinase; CK, casein kinase; CLK, CDC-like Kinase; CSK, C-terminal Src kinase; DAPK, Death-Associated Protein Kinase; DYRK, dual-specificity tyrosine-phosphorylated and regulated kinase; eIF, eukaryotic translation initiation factor; EF2K, elongation-factor-2-kinase; EPH, ephrin; ERK, extracellular-signal-regulated kinase; FGF-R, fibroblast-growth-factor receptor; GCK, germinal centre kinase; GSK, glycogen synthase kinase; HER4, V-erb a erythroblastic leukemia viral oncogene homolog 1; HIPK, homeodomain-interacting protein kinase; IGF, insulin-like growth factor; IKK, inhibitory  $\kappa$ B kinase; IR, insulin receptor; IRAK, Interleukin-1 Receptor-Associated Kinase; IRR, insulin related receptor; JAK, Janus Kinase; JNK, c-Jun N-terminal kinase; Lck, lymphocyte cell-specific protein tyrosine kinase; LKB1, MO25, STRAD, Ser/Thr Kinase 11; MAPKAP-K, MAPK-activated protein kinase; MARK, microtubule-affinity-regulating kinase; MEKK, mitogen-activated protein kinase kinase kinase; MELK, maternal embryonic leucine-zipper kinase; MINK, misshapen-like kinase; MLCK, smooth-muscle myosin light-chain kinase; MLK, mixed lineage kinase; MNK, MAPK-integrating protein kinase; MSK, mitogen- and stress-activated protein kinase; MSPK, Myristoylated and Palmitoylated serine/threonine protein Kinase; MST, mammalian homologue Ste20-like kinase; NEK, NIMA (never in mitosis in *Aspergillus nidulans*)-related kinase; NUA, SnF1-like Kinase; OSR, Oxidative Stress Responsive; PAK, p21-activated protein kinase; PHK, phosphorylase kinase; PDK, 3-phosphoinositide-dependent protein kinase; PIM, provirus integration site for Moloney murine leukaemia virus; PKA, cAMP-dependent protein kinase; PKB, protein kinase B (also called Akt); PKC, protein kinase C; PKD, protein kinase D; PLK, polo-like kinase; PRAK, p38-regulated activated kinase; PRK, protein kinase C-related kinase; RIPK, receptor interacting protein kinase; ROCK, Rho-dependent protein kinase; RSK, p90 ribosomal S6 kinase; S6K, S6 kinase; SGK, serum- and glucocorticoid-induced kinase; Src, sarcoma kinase; SRPK, serine-arginine protein kinase; STK, Serine / Threonine Kinase; SYK, spleen tyrosine kinase; TAK, Transforming growth factor beta activated kinase; TAB, TAK1 binding subunit; TAO, thousand and one amino acid protein kinase; TBK, TANK-binding kinase; TIE, Tunica Internal Endothelial cell kinase; TLK, tousled-like kinase; TrkA, Neurotrophic tyrosine kinase, receptor, type 1 TTK, Phosphotyrosine picked threonine kinase; VEGFR, vascular endothelial growth factor receptor; YES, Yamaguchi sarcoma viral oncogene homolog; ZAP, zeta chain associated protein kinase.

**Figure 3.** The specificities of BMP pathway inhibitors, Dorsomorphin (A) and LDN-193189 (B), against a panel of up to 121 kinases. The results are presented as bars indicating percentage activity remaining for each kinase (averages of two duplicate determinations) in the presence of the indicated concentration of the inhibitor compared with a control lacking

the inhibitor  $\pm$  standard deviation. Further details of the assays are given in the Methods section. The results are ranked according to the percentage activity remaining when the assays were performed in the presence of the indicated inhibitors at 1  $\mu$ M. Protein kinases referred to in the Results section are indicated by red arrows. Abbreviations for protein kinases are described in the Legends to Figure 2.

**Figure 4:** Inhibition of ALK3 and ALK5 by inhibitors of the TGF $\beta$  and BMP pathways. ALK3 (**A**) and ALK5 (**B**) were assayed as described in the Methods section in the presence or absence of the indicated concentrations of TGF $\beta$  pathway inhibitors SB-431542, SB-505124, LY-364947 and A-83-01. The assay samples were resolved by SDS-PAGE, and the gels were Coomassie-stained, dried and analysed by  $^{32}$ P autoradiography. For percentage activity remaining determinations, Coomassie stained bands corresponding to substrate proteins were excised,  $^{32}$ P-incorporation measured and the resulting cpm used as a percentage of control. (**C**) As above, except that ALK3 and ALK5 were assayed in the presence or absence of the indicated concentrations of Dorsomorphin.

**Figure 5.** Inhibition of TGF $\beta$  and BMP pathways by LDN-193189. Serum-starved Human keratinocyte (HaCaT) cells were treated with the indicated concentrations of LDN-193189 for 2h and then treated with BMP-2 (25 ng/ml) (**A**) or TGF $\beta$  (50 pM) (**B**) for 1h. Extracts were resolved by SDS-PAGE and transferred to nitrocellulose membranes, which were analysed by Western Blotting using phospho-Smad1/5/8, Smad1, phospho-Smad2, Smad2/3 and GAPDH antibodies.

**Figure 6.** LDN-193189 inhibits ALK2 and ALK3 in vitro. ALK2 (**A**), ALK3 (**B**), ALK4 (**C**) and ALK5 (**D**) were assayed as described in the Methods section in the absence or presence of the indicated concentrations of LDN-193189. The assay samples were resolved by SDS-PAGE, and the gels were Coomassie-stained, dried and analysed by  $^{32}$ P autoradiography. For IC<sub>50</sub> determinations, Coomassie stained bands corresponding to substrate proteins were excised,  $^{32}$ P-incorporation measured and the resulting cpm plotted against concentrations of LDN-193189 used.

**Figure 7.** Inhibition of NOD-RIPK2 pathways by LDN-193189. RAW 264.7 cells (Mouse leukaemic monocyte macrophage cells) were incubated with indicated concentrations of LDN-193189 for 2h prior to treatment of cells with 15  $\mu$ M Meso-DAP for 1h. Extracts were resolved by SDS-PAGE and transferred to nitrocellulose membranes, which were analysed by Western Blotting using phospho-ERK1/2 and total ERK1/2 antibodies.



**Table 1: Summary of the reported potencies of inhibitors of TGF $\beta$  and BMP pathways.**

**(A)** IC<sub>50</sub> values for the inhibitors of the TGF $\beta$  pathway. The appropriate reference for each determination is indicated. Abbreviations: TGF $\beta$ -induced P-Smad2, TGF $\beta$ -induced phosphorylation of Smad2 in cells as detected by Western blotting; CAGA-Luc, CAGA-Luciferase Reporter Activity dependent on TGF $\beta$  signalling measured in a cell-based assay; ca, constitutively active; nd, not determined. For detailed methodologies, refer to appropriate references indicated **(B)** IC<sub>50</sub> values for the inhibitors of the BMP pathway. The appropriate reference for each determination is indicated. Abbreviations: BMP-induced P-Smad1, BMP-induced phosphorylation of Smad1 in cells as detected by Western blotting; ca, constitutively active. BRE-Luc, BMP-Responsive Luciferase reporter activity measured in a cell-based assay. For detailed protocols, see indicated references.

**A.**

Assay	SB-431542	SB-505124	LY-364947	A-83-01
TGF $\beta$ -induced P-Smad2	0.5-1 $\mu$ M [11]	0.5 $\mu$ M [16]	0.135 $\mu$ M [19]	nd
ALK4 in vitro kinase assay	0.14 $\mu$ M [15]	0.129 $\mu$ M [16]	nd	nd
ALK5 in vitro kinase assay	0.094 $\mu$ M [11]	0.047 $\mu$ M [16]	0.058 $\mu$ M [19]	nd
CAGA-Luc (Cells)	0.25 $\mu$ M [16]	0.1 $\mu$ M [16]	nd	nd
caALK4 (CAGA-Luc)	0.75 $\mu$ M [11]	nd	nd	0.100 $\mu$ M [20]
caALK5 (CAGA-Luc)	0.5 $\mu$ M [11]	nd	nd	0.012 $\mu$ M [20]
caALK7 (CAGA-Luc)	1-2 $\mu$ M [11]	nd	nd	0.030 $\mu$ M [20]

**B.**

Assay	Dorsomorphin	LDN-193189
BMP-induced P-Smad1/5/8	0.47 $\mu$ M [24]	0.005 $\mu$ M [25]
caALK2 (BRE-Luc)	0.20 $\mu$ M [24]	0.005 $\mu$ M [6]
caALK3 (BRE-Luc)	0.50 $\mu$ M [24]	0.03 $\mu$ M [6]
caALK6 (BRE-Luc)	5-10 $\mu$ M [24]	0.15 $\mu$ M [6]

**Table 2: Potencies of compounds developed as TGF $\beta$  and BMP pathway inhibitors against some other kinases.** IC<sub>50</sub> values were determined from multiple assays carried out at ten different inhibitor concentrations. Abbreviations for protein kinases are described in the Legends to Figure 2.

Kinase	SB-431542	SB-505124	LY-364947	A-83-01	LDN-193189
RIPK2	0.41 $\mu$ M	0.35 $\mu$ M	0.11 $\mu$ M	0.10 $\mu$ M	0.025 $\mu$ M
CK1 $\alpha$	1.34 $\mu$ M	19.44 $\mu$ M	2.27 $\mu$ M	15.66 $\mu$ M	3.61 $\mu$ M
CK1 $\delta$	0.92 $\mu$ M	3.38 $\mu$ M	0.22 $\mu$ M	3.42 $\mu$ M	0.92 $\mu$ M
CK1 $\epsilon$	0.38 $\mu$ M	1.60 $\mu$ M	1.34 $\mu$ M	4.59 $\mu$ M	14.24 $\mu$ M
CK1 $\gamma$	>100 $\mu$ M	>100 $\mu$ M	43.97 $\mu$ M	28.55 $\mu$ M	98.92 $\mu$ M
GCK	>100 $\mu$ M	>100 $\mu$ M	7.91 $\mu$ M	2.22 $\mu$ M	0.08 $\mu$ M

Figure 1




Figure 1

Figure 2AB





Figure 2

Figure 2CD





Figure 2

Figure 3




Figure 3

Figure 4




Figure 4

Figure 5





Figure 5



**Figure 6**



**Figure 6**

Figure 7




Figure 7